

Tamias striatus. By Dana P. Snyder

Published 25 May 1982 by The American Society of Mammalogists

Tamias Illiger, 1811

Tamias Illiger, 1811:83. Type species *Sciurus striatus* Linnaeus, 1758:64, by monotypy.

CONTEXT AND CONTENT. Order Rodentia, Family Sciuridae, Subfamily Sciurinae. There is disagreement over whether all chipmunks (*Tamias*, *Eutamias*) should be included in *Tamias*, or only the eastern chipmunk on North America. See Remarks.

Tamias striatus (Linnaeus, 1758)

Eastern Chipmunk

[*Sciurus*] *striatus* Linnaeus, 1758:64. Type locality upper Savannah River, South Carolina; fixed by Howell, 1929:14.

Tamias americana Kuhl, 1820:69. Based on material in Bullock's museum; locality given only as "In America septentrionali."

Sciurus (Tamias) lysteri Richardson, 1829:181. Type locality Penetanguishene, Ontario.

Tamias striatus, Baird, 1857:55. First use of name combination.

CONTEXT AND CONTENT. Context in generic account above. In his 1929 revision Howell recognized five subspecies and six more have been described since (Hall, 1981). Ellis (1979), basing his conclusions on cranial and external morphology and to some extent on coloration, accepted only five subspecies. Because Ellis' revision is not yet widely available, the eleven previously described forms are listed below (see Distribution).

T. s. doorsiensis Long, 1971:201. Type locality Peninsula State Park, Door Co., Wisconsin.

T. s. fisheri Howell, 1925:51. Type locality Merritt's Corners, 4 mi E Ossining, Westchester Co., New York (see Paradiso, 1963).

T. s. griseus Mearns, 1891:231. Type locality Fort Snelling, Minnesota.

T. s. lysteri (Richardson, 1829:181), see above. Regarding authorship see Merriam (1886).

T. s. ohionensis Bole and Moulthrop, 1942:135. Type locality Cincinnati, Hamilton Co., Ohio.

T. s. peninsulae Hooper, 1942:1. Type locality Barnhart Lake, 3 mi SE Millersburg, Presque Isle Co., Michigan.

T. s. pipilans Lowery, 1943:235. Type locality 5 mi S Tunica, West Feliciana Parish, Louisiana.

T. s. quebecensis Cameron, 1950:347. Type locality St. Félicien, Lake Saint John Co., Quebec.

T. s. rufescens Bole and Moulthrop, 1942:130. Type locality Chesterland Caves, Chester Township, Geauga Co., Ohio.

T. s. striatus (Linnaeus, 1758:64), see above (*americana* Kuhl a synonym).

T. s. venustus Bangs, 1896:137. Type locality "Stilwell, I. T.," now Stilwell, Adair Co., Oklahoma.

DIAGNOSIS. *Tamias striatus* differs from other chipmunks (subgenus *Eutamias*, subgenus *Neotamias*) in the absence of P3, lack of fusion of hypohyal and ceratohyal bones, and presence of keel on ventral tip of baculum. It differs from these subgenera also in having a relatively shorter tail (less than 38% of total length versus more than 40% of total length in others). The innermost paired dorsal stripes are about twice the width of other stripes in *T. striatus* rather than being about equal in width to the remaining stripes as in the other chipmunks. The eastern chipmunk differs from small terrestrial squirrels in the form of the infraorbital foramen which consists of a short oval passageway perforating the zygomatic plate rather than a canal opening forward of the plate (White, 1953). Additional diagnostic characters are given by Bryant (1945), Howell (1938), and White (1953).

GENERAL CHARACTERS. The eastern chipmunk is a small, moderately heavy-set, sciurid with prominent longitudinal stripes (Fig. 1). Mid-dorsally a dark brown, quite narrow stripe extends from back of head to rump; this stripe is bordered on either side by a strip, about twice as wide, of agouti hairs varying from a grayish to reddish-orange or brownish in appearance. Lateral to these are two narrow, dark-brown stripes enclosing a slightly wider, yellowish-white stripe. These stripes fade into the yellowish- or reddish-brown rump. Yellow-orange sides grade into the whitish underside. The prominent eyes are bordered above and below by light facial stripes which are in turn bordered by darker stripes. The tail is dorsoventrally flattened, well haired but not bushy, and dark above with light gray border. The underside of the tail has a central reddish-orange strip enclosed by narrower blackish strips and is edged with light gray. Overall appearance is of a grayish- to reddish-brown small mammal. Intensity and shade of coloration varies geographically; the palest forms occur in southern Ontario, dark red forms in parts of the southern Appalachians, and the most brightly colored individuals in southwestern portions of the range. Along the northern parts of the range, reddish effects are reduced, giving a darker gray, less colorful, appearance. Ears are fairly prominent, less than 20 mm to tip from notch, and more or less rounded at apex. There are four functional toes on the forefeet (pollex vestigial with a flat nail) and five on the hindfeet. Internal cheek pouches are large. There are four pairs of mammae—1 pectoral, 2 abdominal, and 1 inguinal. Smallest forms occur in the southeastern portion of the range and largest in the northwest and southwest. Average measurements (mm) for seven populations from widely scattered points over the geographic range of the species are: total length, 225 to 268; length of tail, 72 to 101; greatest length of skull, 39.4 to 43.8. Adult weights range from less than 80 g to 125 g or more.

The skull (Fig. 2) is lightly built and somewhat elongate with relatively small and weak postorbital processes. The zygomatic plate slants upward at approximately 45° from the occlusal plane of the cheekteeth. Small masseteric tubercles lie ventral and partially mesial to the infraorbital foramen. The frontoparietal suture is not ankylosed and temporal ridges do not meet to form a sagittal crest. There is no temporal foramen in the squamosoparietal suture. The superior outline of skull is shallowly convex. The braincase is shallow and the lambdoid crest is well developed. Head of the malleus is more elongate than in other chipmunks and the auditory bulla has two transbullar septa. Dental formula is $i\ 1/1$, $c\ 0/0$, $p\ 1/1$, $m\ 3/3$, total 20. A succinct characterization of all the chipmunks as a group (including the subgenera *Eutamias* and *Neotamias*) was given by Moore (1959).

DISTRIBUTION. The eastern chipmunk ranges from the vicinity of Lake Manitoba eastward to the Quebec coastline just north of Anticosti Island and southward to the Gulf states (Fig. 3). It is present on Cape Breton Island but absent from Anticosti Island and (originally) from Newfoundland where, in the 1960's, chipmunks from Nova Scotia were introduced at three sites (Northcott et al., 1974). On the west, discontinuous outposts occur in wooded habitats quite far from typical deciduous forest regions, particularly in river valleys (Black et al., 1977; Howell, 1929). In Louisiana it occurs only in a few parishes near the Mississippi River (Lowery, 1974), and in Florida only in the north-west part of the state near the Alabama line (Stevenson, 1962). It is absent from much of the coastal plain and some adjacent areas of piedmont in the Carolinas and Georgia. *T. striatus* occurs near human habitations as well as in uninhabited areas and is an "urban" as well as a wilderness species.

FOSSIL RECORD. Chipmunks (sensu lato) appear to have been a distinct stock since at least the late Oligocene (Black, 1963) and are represented by later Tertiary fossil material from several widely scattered sites in North America (Black, 1963; Klingener, 1968; Lindsay, 1972; Sutton, 1972). Place of origin of



FIGURE 1. Photograph of *Tamias striatus* from Allegheny National Forest, Pennsylvania, taken by Kenneth E. Fisher.

this group and its later Tertiary migration history is not known, but several scenarios have been suggested (Black, 1972; Ellis and Maxson, 1979; Moore, 1961; Nadler et al., 1977). Ellis and Maxson (1979) postulated the time of divergence of chipmunks from the tree squirrel lineage as 38 million years BP (Oligocene) and that *Tamias* (sensu stricto) diverged from the other American chipmunks during late Miocene (10 million years BP).

Several Pleistocene species closely related to the eastern chipmunk have been described, but only one, *T. aristus* (Ray, 1965), appears to be distinct from *T. striatus*. Specimens definitely referable to *T. striatus* are common in late Pleistocene and early Recent deposits, particularly in caves, sinks, and fissures, and around prehistoric Indian sites within the current range of the species (Guilday, 1963; Ray, 1965). The stratigraphic range extends from late Sangamonian to Recent (Kurtén and Anderson, 1980). In late Pleistocene to early Recent time (approximately 8,000 to 11,000 years BP) the geographic range of *T. striatus* was at least several hundred km west of its present known limits (Dalquest et al., 1969; Roth, 1972). Ray (1965) summarized the Pleistocene to Recent material.

FORM. Pelage pattern and, more strikingly, color vary geographically (Allen, 1938). Neither melanistic nor albino variants are common, but there are records of both from central and north-eastern parts of the range (Goodpaster and Hoffmeister, 1968; D. Smith and L. Smith, 1972). There are two molting peaks each year, but there is difference of opinion over whether the same chipmunk molts twice in 1 year (Bole and Moulthrop, 1942; Ellis, 1979; Howell, 1929; Yerger, 1955). Yahner et al. (1979) found numerous sebaceous glands associated "... with hair follicles around the anal orifice and in the entire perineal region ...". Aggregations of glandular tissue, which they speculated might be a source of reproductive pheromones, were also noted in the walls of the anal canal extending into the perineum. Quay (1965) found sebaceous and mucous glands but not sudoriferous glands in the oral lips and angle of *T. striatus*.

Comparative descriptions of the baculum have been published by Burt (1960) and Wade and Gilbert (1940). Average length of 31 bacula, mostly from Michigan, was 4.15 mm (Burt, 1960). The length of the os clitoridis is approximately 60% that of the baculum (Layne, 1954). The cutaneous maximus muscle is poorly developed with only a dorsal and a pudendal portion (Woods and Howland, 1977). Many detailed structural characteristics of the skeleton and musculature are given in the comparative descriptions by Bryant (1945).




FIGURE 2. Dorsal, ventral, and lateral views of skull and lateral view of mandible of male *Tamias striatus* from West Feliciana Parish, Louisiana (Louisiana State University Museum of Zoology No. 16177). Drawing by H. D. Pratt from Lowery (1974), *The mammals of Louisiana and its adjacent waters*, by permission of Louisiana State University Press. Cranium measures approximately 44 mm.

Neuroanatomy has been little studied, but Simpson (1914) found a complete decussation of the pyramidal tracts which, after crossover, travel in the dorsal columns of the spinal cord as far as the lower sacral vertebrae, with a considerable number of fibers disappearing in the cervical region. Sealander (1964) gave values for several blood parameters. Anthony et al. (1963) described the histology of the digestive organs; they found no difference between active and hibernating chipmunks or between sexually active and sexually quiescent individuals. Mossman et al. (1932) found that *Tamias striatus* had large seminal vesicles, large prostate and Cowper's glands, and a small bulbar gland with no intermediate portion. The testes of an old adult in breeding condition measured $10 \times 10 \times 17$ mm. Dudkiewicz (1968) described the histology of the female reproductive tract. The ovary is approximately 2.5 mm long by 1.5 wide. The mature graafian follicle attains a diameter of about 560μ , and the corpora lutea of pregnancy are approximately 800μ in diameter.

FUNCTION. Mean minimum heart rate of normothermic animals at ambient temperatures of 30° to 31°C was found to be 165 beats/min. Rates increased at higher and at lower temperatures, reaching 530 at -5°C (Wang and Hudson, 1971). Spontaneous cardiac activity can be maintained in isolated atria at temperatures as low as 0° to 4°C ; activity of isolated ventricles ceases at a slightly higher temperature (Angelakos et al., 1969; Lyman and Blinks, 1959). At about 20°C captive animals showed annual cycles in erythrocyte count (higher in summer than winter) and blood sugar (increased from January to July, then decreased) (Woodward and Condren, 1945). Wang and Hudson (1971) found that mean oxygen consumption at ambient temperatures from

28.5° to 32°C was $1,030 \text{ mm}^3 \text{ g}^{-1} \text{ h}^{-1}$. Body temperatures were between 36° and 40.3°C at ambient temperatures between -7° and +32°C. Above 32°C body temperature climbed and above 36°C chipmunks had difficulty dissipating the heat load. Thermoneutrality was between 28.5° and 32°C. However, Neumann (1967) found the thermoneutral zone to extend from 28° to 36°C, and within this range mean body temperature was 39.2°C and mean oxygen consumption $1,250 \text{ mm}^3 \text{ g}^{-1} \text{ h}^{-1}$.

Neff (1968) measured gross metabolic response in active chipmunks exposed to $3^\circ \pm 2^\circ \text{C}$ for 30 days; body temperatures averaged $37^\circ \pm 1^\circ \text{C}$, the same as controls kept at $24^\circ \pm 1^\circ \text{C}$. He concluded that chipmunks have a well-developed capacity for acclimation to reduced temperatures by gross metabolic changes similar to those observed in homeotherms such as laboratory rats. Neal and Lustick (1975) found that artificial infra-red radiation at energy levels which might be expected in nature caused a downward shift of 3.5° to 8.2°C in the thermoneutral zone. This reduced O_2 consumption, indicating that basking might be important to chipmunks. Importance of the nasal passage in heat exchange was demonstrated by Schmid (1976) who found a gradient of 10.7°C between back of passage (cribriform plate) and exhalent air. Chipmunks respond to hypoxia (8% O_2) by an elevation in heart and respiratory rate and decrease in body temperature (Faleschini and Whitten, 1975). Blake (1977) found that minimum water requirements of captive chipmunks varied from about 24% of their weight per day in the reproductive season to 4% to 5% of weight during the hibernating season. Captive chipmunks studied by Forbes (1967) with a mean weight of 115 g utilized an average of 16.4 ml of H_2O per day.

From late fall to early spring chipmunks are mostly underground in various degrees of torpor, but sometimes appear above ground in favorable weather. Some remain dormant throughout the winter. Juveniles tend to remain active until later in the fall than do adults. In captivity a variable proportion of chipmunks become torpid at ambient temperatures ranging from 0° to 22°C, but only during the winter and spring months (Pivorun, 1976b; Wang and Hudson, 1971). The annual torpidity cycle appears to be endogenous and is probably independent of temperature (Scott and Fisher, 1972). However, Woodward and Condren (1945) reported that if blood sugar of captive chipmunks were lowered by insulin, torpidity could be easily induced by a temperature of 5° to 10°C. Differences in photoperiod do not seem to affect entry into torpor but do affect length of torpor (Brenner and Lyle, 1975). Body temperature usually remains several degrees (often more than 10) above ambient, but two individuals observed by Wang and Hudson (1971) maintained only a 1° gradient during torpor at 4°C ambient. Torpidity allows chipmunks to reduce their metabolic needs by 85% or more (Neumann, 1967; Pivorun, 1976a; Wang and Hudson, 1971). Length of torpid bouts varies from less than 24 hours to several days and is a function, according to Pivorun (1977), of body temperature attained during torpor. Winter inactivity occurs even in northern Florida (Stevenson, 1962) and continues as long in some southern areas as in the north (Engels, 1951). Arousal occurs fairly rapidly, taking from less than 0.5 h to 2 h depending on depth of torpor. Temperatures in abdominal areas tend to lag behind those of the thorax and other anterior regions of body during warming (Schaffer and Bullard, 1970; Wang and Hudson, 1971). Schaffer (1971) discussed the importance of brown fat in arousal from torpidity. Anthony et al. (1967) and Munro and Anthony (1965) discussed other physiological aspects of hibernation. Captives exhibit an annual cycle in body weight (Allen, 1938; Brenner and Lyle, 1975; Panuska, 1959; Schaffer, 1973), but Forbes (1966a) found no evidence of fall accumulation of fat in free-ranging chipmunks.

Thyroid activity appears to be greatest in the fall and lowest in summer, and females show higher activity than males (Fisher, 1972; Kodrich and Tryon, 1971). Adrenal weights of captive chipmunks at room temperature increased nearly 3-fold from a January low to a plateau from April to September and then regressed (Woodward and Condren, 1945). Clulow et al. (1969) found that adrenocortical activity was positively correlated with density in populations of wild caught animals introduced onto islands.

ONTOGENY AND REPRODUCTION. Most matings occur between late February and early April and again from late June to early July. Males come into breeding condition in mid-February or earlier and testes begin to regress in July or August. Females are in breeding condition by the time of emergence, usually in March. Females may breed in either the spring or summer period or both (L. Smith and D. Smith, 1972; Yerger, 1955). D. Smith and L. Smith (1975) found that estrus, as defined by morphological characteristics of the external genitalia, lasted




FIGURE 3. Geographic range of *Tamias striatus* showing the 11 described forms and the subspecies considered valid by Ellis (1979). Named forms, bounded by dashed lines are: 1, *T. s. doerriensis*; 2, *T. s. fisheri*; 3, *T. s. griseus*; 4, *T. s. lysteri*; 5, *T. s. ohionensis*; 6, *T. s. peninsulae*; 7, *T. s. pipilans*; 8, *T. s. quebecensis*; 9, *T. s. rufescens*; 10, *T. s. striatus*; 11, *T. s. venustus*. Subspecies according to Ellis, denoted by shading are: A, *T. s. griseus*; B, *T. s. lysteri*; C, *T. s. pipilans*; D, *T. s. striatus*; E, *T. s. venustus*. Populations introduced into Newfoundland are indicated by solid circles. Adapted from Hall (1981) and Ellis (1979). Drawn by K. A. Sargent.

at least 3 days and up to 10 days. However, on behavioral criteria Elliott (1978) considered that estrus lasted only about 6.5 h. Gestation is 31 to 32 days (Allen, 1938; Burt, 1940). Behavior of a captive animal during parturition was described by Allen (1938). Litter size averages between 4 and 5. Newborn weigh 2.5 to 5.0 g; a female measured by Allen (1938) was 66 mm in total length, with a 12 mm tail, and a 7.5 mm hindfoot. By 1 month of age juveniles are 140 mm long and weigh 30 g; eyes are open, body is well furred, and they have an adult appearance. They emerge at 5 to 7 weeks of age (Pidduck and Falls, 1973; L. Smith and D. Smith, 1972; Yerger, 1955). Adult size and permanent dentition are achieved by 3 months. By 1 year the teeth are well worn. Additional growth-rate data were given by Pidduck and Falls (1973) and Yahner (1978a). Individuals are usually not sexually mature until after their first winter, but some spring-born females mate in the following summer. Males rarely, if ever, become sexually mature their first summer (L. Smith and D. Smith, 1972; Yerger, 1955). Seasonal changes in the male reproductive tract were studied by Neff and Anthony (1963). Characteristics of teeth, skull, and long bones have been used to determine age (Allen, 1938; Ellis, 1979; Yerger, 1955).

ECOLOGY. *Tamias striatus* inhabits primarily deciduous wooded areas; it occurs in open bushy habitats as well as mature forests, especially in habitats with abundant crevices for refuge along with elevated observation and vocalization posts. Ryan and Larson (1976) reported habitat characteristics of chipmunks in residential areas. Density is temporally and geographically variable, ranging from 0.3 to 37.6 per ha (Yerger, 1953). Annual variation is commonly 2- to 5-fold and tends to be greater in areas with the lower population densities. Sex ratios of trapped chipmunks are quite variable; differences likely result from chance variation in small samples and from differences between sexes in sizes of home ranges. Tryon and Snyder (1973a) published life tables for populations in Pennsylvania and Vermont. Male-female ratios of 1,507 juveniles from several different locations varied from 52:48 to 45:55, but the overall ratio was 50:50. Survival rates were fairly constant over much of the adult life but varied geographically and by season of birth; spring litters survived better than those born in late summer. Some chipmunks lived at least into year 8; average annual survival and mean life

expectancy from time of first capture (shortly after emergence) was 0.43 and 1.3, respectively. There is annual variation in age structure as a result of different numbers of juveniles entering the population in different years. Roberts (1976) made an extensive analysis of age, sex, and native/immigrant population structures.

Panuska and Wade (1956), Thomas (1974), and Elliott (1978) made detailed studies of burrow systems. Some are single tunnels, but more complex systems contain one or more large storage chambers and various smaller galleries and short pockets in which food or debris is stored. Length is up to 10 m, depth less than 1 m. The nest, of chewed or crushed leaves, occupies one of the larger chambers. A female in Michigan utilized a hollow tree as a nest site in which to bear and raise a litter (Getty, 1979). Entrances are 40 to 60 mm in diameter; usually only one is in active use, but plugged alternate passages frequently reach the surface.

Mean minimum area home range varies temporally and geographically from 0.03 to 0.40 ha with individual home ranges varying from under 100 m² to more than a hectare. Elliott (1978) found that ranges usually were more or less symmetrical with centrally located burrows and that use-intensity was highest near the burrow and gradually decreased toward the periphery. Detailed breakdowns of ranges according to age, sex, time of year, and other factors were made by Ickes (1974) and West (1975). Chipmunks make long excursions outside their usual range, in part to exploit concentrated food sources. Roberts (1976) found that some resident (non-dispersing) individuals have lifetime range lengths of at least 484 m; dispersal movements extended at least 857 m, with males making longer movements. Mares et al. (1976) were able to induce a contraction in size of ranges (from an average size of 0.02 to 0.11 ha) by providing a concentrated and unlimited food source within core areas. Most chipmunks are relatively sedentary and retain their burrow systems and range for long periods, often for life (Yahner, 1978b).

Various aspects of the diet and feeding ecology have been studied by Elliott (1978), Forbes (1966b), Frost (1968), Graybill (1970), Wrazen and Svendsen (1978), and Yerger (1955). Graybill (1970) listed published references to food items. Staple, over-wintering food consists of seeds, nuts, and acorns larder-hoarded in burrow systems. Frost (1968) and Wrazen and Svendsen (1978) found that nuts and acorns, apparently from the winter food cache, were used during the spring and summer. Elliott (1978) estimated that some hoards which he excavated in May and June could have lasted their owners well into the next winter. Yahner (1975) and Elliott (1978) discussed the significance of scatter-hoarding outside burrows. Invertebrates (especially insects) and fungi also are important foods during the growing season (Elliott, 1978; Frost, 1968; Wrazen and Svendsen, 1978). Chipmunks prey on frogs, snakes, birds, and small mammals (see Graybill, 1970). Graybill (1970) found that captive chipmunks kept under outdoor conditions in Pennsylvania utilized an average of 32.7 to 35.7 kcal per individual per day during a 1 year period. He estimated that free-ranging chipmunks in areas he studied consumed more than half of their potential food resource.

Tamias striatus is prey for snakes, hawks, weasels, foxes, and bobcats (Hamilton and Whitaker, 1979). Peifer (1979) saw a great blue heron, *Ardea herodias*, take chipmunks, and Symington (1907) observed a "large frog" capture and eat a chipmunk.

Antibodies to the La Crosse strain of California Encephalitis virus have been detected in chipmunks in Ohio and Wisconsin, and the virus has been isolated from Wisconsin chipmunks (Berry et al., 1975). Known protozoan parasites are *Trypanosoma tami-* asi, *Eimeria ovata*, and *E. vilasi* (Dorney, 1966; Duncan, 1968; Riedel, 1975). Helminths reported are: Cestoda—*Hymenolepis diminuta*, *H. sp.*, *Taenia mustelae* (bladderworm), *Cladotenia globifera* (plerocercoid), and *Paruterina rauschi* (plerocercoid) (Freeman, 1956, 1957, 1959; Rausch and Tiner, 1948); Trematoda—*Scaphistomum pancreaticum* and *Postharmostomum helici* (Elwell, 1967; Rausch and Tiner, 1948); Nematoda—*Trichostrongylus sp.*, a metastrongyloid (paratenic), *Rictularia halli*, *Spirura michiganensis*, *Trichuris madisonensis*, and *Capillaria tamias-striati* (Lankester and Anderson, 1966; Rausch and Tiner, 1948; Read, 1949; Tiner, 1950); and *Acanthocephala*—*Macracanthorhynchus hirudinaceus* and *Moniliformis clarki* (Van Cleave, 1953). Infestations by larval *Cuterebra emasculator* are common from July to October, generally occurring near the urogenital openings or between them and base of tail. Infested animals may exhibit splenomegaly, thymic involution, anemia, elevated leucocyte counts, and purulent wounds after larval emergence. There is little direct evidence of death from this parasitism (Bennett, 1973; Dorney, 1965; McKinney and Christian,

1970). *Aedes cinereus* and *A. triseriatus* commonly feed on *T. striatus* (Wright and DeFoliart, 1970). Ticks and mites found on chipmunks include *Dermacentor variabilis*, *Haemaphysalis leporispalustris*, *Ixodes marxi*, *I. muris*, *I. dammini*, *Haemolaelaps glasgowi*, *Eulaelaps stabularis*, *Euschongastia peromysci*, *Sciurocoptes tami-* as, and *Trombicula harperi* (Carey et al., 1980; Fain and Hyland, 1970; Jones and Thomas, 1980; Parsons, 1962). Hypopial nymphs recorded from *T. striatus* are *Aplodontopus sciuricola* and one attributed to *Dermacarus hylandi* (Fain et al., 1971). Seasonal infestation rates by *A. sciuricola* were discussed by Tadmowski and Hyland (1979) who found 95% of 77 specimens infested. Fleas recorded from *T. striatus* are *Tamiothrips grandis*, *Stenoponia americana*, *Megabothris acerbus*, *M. quirini*, *Ceratophyllus gallinae*, *Ctenophthalmus pseudagrytes*, *Monopsyllus vison*, *Orchopeas howardii*, *O. caedens*, *O. leucopus*, *Nosopsyllus fasciatus*, *Epitedia wenmanni*, and *Peromyscopsylla catatula* (Amin, 1976; Lovejoy and Gaughan, 1975; Parsons, 1962). Parsons (1962) listed lice found on eastern chipmunks as *Hoplopleura erratica* and *Enderleinellus tami-* asus.

Aerial spraying of forests in Ohio with 0.81 kg/ha of malathion resulted in a 55% reduction in the chipmunk population (Giles, 1970). Chronic exposure of free-ranging chipmunks to gamma radiation resulted in a temporary weight loss of several grams (Buech, 1977). Acute exposure of free-ranging individuals to gamma radiation (200 to 400 roentgens) resulted in an increase in survival rate (Tryon and Snyder, 1973b), and in a reduction of range size and/or mobility of a population normally exhibiting relatively large home ranges (Snyder et al., 1976).

BEHAVIOR. Social organization was studied by Elliott (1978), Ickes (1974), and Yahner (1978c, 1978d). More restricted aspects of social behavior were reported by Aniskowicz and Vailancourt (1979), Brenner et al. (1978), Dunford (1970), Herbig (1978), Wolfe (1966), Yerger (1953), and others. A solitary individual occupying a single burrow system is the basic social unit, except for the brief mother-offspring grouping (Yahner, 1978c). Individuals occupy broadly overlapping home ranges, but each spends most of its time within a core area defended by threat behavior, or by a chase which is usually short (3 to 4 m), but may extend up to 15 m (Ickes, 1974; Yahner, 1978c). Reversal of the dominant-subordinate relationship usually occurs if the chase extends into the core area of the intruder. Most chases do not result in physical contact, but biting by the dominant, and/or actual fights occur occasionally though no serious injury seems to result (Dunford, 1970; Elliott, 1978; Ickes, 1974). Space-independent factors affecting dominance relationships are weight, sex, age, and previous encounter experience (Brenner et al., 1978; Ickes, 1974; Wolfe, 1966). When they meet on "neutral" ground, chipmunks tolerate conspecifics at distances of 0.5 to 1.0 m, which are maintained by periodic short chases (Elliott, 1978; Yahner, 1978c). Yahner (1978d) found that behavior is more stereotyped in adults than young, in males than females, in breeding than nonbreeding seasons, away from rather than near home sites, and when a conspecific is nearby rather than when no conspecific is in the vicinity. Agonistic behavior accounted for less than 3% of a chipmunk's time budget.

During breeding periods, males are active early in the morning, inspecting burrow entrances, core areas of other chipmunks, and conspecifics encountered, and congregating on ranges of estrus females. According to Yahner (1978c), when only two or three males were involved in a mating bout, the dominant always copulated with the female and prevented access by others. If a large number of males (7 to 8) were present, however, the dominant male spent so much time chasing rivals that he never copulated with the female. One male may copulate several times with the same female and different males may also copulate with her during her estrus period.

Some mothers transport young to a neighboring burrow prior to emergence. Others may make a separate nest in their own burrow. Mothers become intolerant a few days after young emerge and the latter establish their own home within 2 weeks or less after emergence. Until then, they stay close to their burrows, remain friendly to each other, and frequently group together (Elliott, 1978; Yahner, 1978c). Roberts (1976) believed that most young make only short dispersal movements and that the average movement of a juvenile from its natal burrow to its later residence is less than the average radius of home range.

Captives studied by Wolfe (1969) explored extensively when first placed in an unfamiliar cage, but this tendency decreased as familiarity with surroundings increased. This behavior first appeared at 26 days of age, 4 days after young gained ability to walk and about a week before their eyes opened. Wolfe (1969)

found that both captive and free-ranging chipmunks generally exhibited neophilia in response to a new object although some transitory neophobia was noted in captives. Herbig (1978) found less neophilia in captives than did Wolfe; she also found that when chipmunks were allowed access to a large unfamiliar enclosure from an adjacent nest box, they more commonly explored visually by sitting at the entrance and peering than by actually entering the area.

Both sexes vocalize prominently; frequencies range from 1 to 11 kHz. Although the sounds form a graded series, at least four types can be distinguished: a single high-pitched "chip," a chip-trill, a series of chips close together referred to as chipping, and a low-pitched "cuk" given repeatedly. The chips are given as frequently as 130 times a minute and for as long as 11 minutes (Seton, 1929); sometimes several chipmunks form a chorus of chipping calls. See Dunford (1970) and Elliott (1978) for sonograms. All calls function as alarm signals, but most especially the chip-trill which is given frequently during an escape rush. Vocalizations also may have a territorial function (Dunford, 1970) or may alert individuals to presence of a predator (Elliott, 1978). There is some decrease in activity in late summer following the second breeding period and prior to juvenile emergence. Yahner (1977) also noted an activity lull after the late winter breeding.

The eastern chipmunk is strictly diurnal, and intensity of activity varies throughout the day; several authors have noted mid-morning and mid-afternoon peaks (Burt, 1940; Dunford, 1972; Wolfe, 1966). However, Yahner (1977) found that no individuals showed a bimodal pattern, but males were more active in morning and less active in afternoon than females. Thibault's (1969) results showed a primarily unimodal pattern with greatest activity between 0900 and 1300 and a peak from 1100 to 1200. This tended to decrease possible conflict with *Tamiasciurus* whose activity peak was between 1600 and 2000. Greatest activity occurs on bright, warm, quiet days (Forbes, 1966b). Graefe (1961) found that 35 to 45 Lux were sufficient to stimulate chipmunks to activity. *Tamias* has an endogenous free-running circadian rhythm of activity and can be entrained to a wide range of schedules (6L:18D to 21L:3D) (DeCoursey, 1972). In activity wheels, chipmunks show a strong directional tendency and apparently get their bearings from the light source or features of the environment (Kavanau, 1969). He raised the possibility that chipmunks may, therefore, use celestial objects and landmarks as navigational aids.

Several investigators have studied homing behavior (Layne, 1957; Seidel, 1961). Individuals released at 100 or 200 m from point of capture were recaptured frequently at or near that point within a day or two, but some were recaptured in the opposite direction (Layne, 1957) or not seen again. Return frequency decreased when release distances were 300 m or more, but Seidel (1961) recovered two of four chipmunks from a distance of nearly 500 m. Return rate of juveniles is much poorer than that of adults. Although these return distances are greater than the span of the usual home range, Roberts' (1976) results demonstrating large lifetime home range size indicate that one cannot be sure the homing chipmunks were released into completely unfamiliar areas. Chipmunks can leap 380 to 610 mm on the level and more than 760 mm from a log or similar elevation (Jackson, 1961). Layne and Benton (1954) determined the average maximum running speed as 11.2 ft/s (12.2 km/h) with an extreme of 15.6 ft/s (17.1 km/h).

GENETICS. Karyotypes derived from bone marrow chromosomes have been discussed and figured (Nadler et al., 1977; Thompson, 1971). The diploid number is 38 and little or no intra-specific karyotypic variation is present. There are four pairs of large metacentric, six pairs of large submetacentric, and one pair of small metacentric autosomes. In addition, 3 pairs of autosomes are large acrocentrics and 4 pairs are small with terminal or sub-terminal centromeres. The X chromosome is a large submetacentric and the Y is minute and dot-like or submetacentric. Fundamental number is 58 (Nadler and Hoffmann, 1970). Based on size and G-bands, nine structural chromosomal differences between *Tamias* (*Tamias*) and *Eutamias* (*Neotamias*) have been demonstrated. *Eutamias* (*Eutamias*), for which G-banding data are not yet available, differs from *Tamias* (*Tamias*) in three autosomal arrangements and in having a larger Y chromosome with some structural differences.

Santiago and Rake (1973) estimated from reassociation kinetics that *Tamias* chromosomes have three different DNA fractions: a unique portion making up 75% of the total; a highly repeated portion (~6,730 copies) constituting 12%; and a section with intermediate repetitiveness (~99 copies) totaling 13%. They also estimated a genome size of 1.3×10^9 nucleotide pairs.

REMARKS. The genus *Tamias* was erected by Illiger (1811) for the chipmunks, those members of his "squirrel" family possessing cheek pouches. Trouessart (1880) created the subgenus *Eutamias* for the Siberian chipmunk and all American forms (including some ground squirrels) except the eastern chipmunk. Merriam (1897) raised *Eutamias* to generic rank, thus leaving the eastern chipmunk as the only species in the genus *Tamias*. Workers on the Siberian chipmunk tended to follow this separation and placed them in *Eutamias* also. Ellerman (1940) and Bryant (1945) argued for a single genus for all chipmunks, but White (1953) marshalled considerable evidence for the generic separation of the eastern chipmunk from all others as proposed by Merriam (1897), and until recently this arrangement was fairly widely followed. However, Nadler et al. (1977) suggested, primarily on the basis of karyotypic comparisons, that all chipmunks should be placed in the genus *Tamias*. Hight et al. (1974), using immunological evidence, seemed to lean toward this view also; but Ellis and Maxson (1979), utilizing a micro-complement fixation technique to quantify immunological characteristics of serum albumen, believed that two genera should continue to be recognized. Regardless of these questions the Recent species can be placed readily into three groups: subgenus *Eutamias*, the Siberian chipmunk, *Eutamias sibiricus*; subgenus *Neotamias*, the western (and northern) North American chipmunks; and subgenus *Tamias*, the eastern North American species. However, firm generic allocation of these groups is premature, and established usage (Hall, 1981) should be retained until additional cytological and biochemical data can be obtained from the Siberian chipmunk for comparison with those already available from the American forms.

Linnaeus' (1754) description of *T. striatus* was based on an American specimen (*vide* Baird, 1858) and the accounts of Catesby (1743) and Edwards (1751). In the 10th edition of *Systema Naturae*, Linnaeus (1758) referred to his 1754 description and cited Kalm (1756) in addition to Catesby and Edwards. The type locality restriction by Howell was based on the region Catesby was known to have traveled. There has never been any confusion concerning the distinctness of *T. striatus* from other North American chipmunks since their discovery, although it was regarded as conspecific with the Siberian chipmunk by some early workers (e.g. Gmelin, 1788; Linnaeus, 1766), who had no knowledge of other New World chipmunks.

LITERATURE CITED

- Allen, E. G. 1938. The habits and life history of the eastern chipmunk, *Tamias striatus lysteri*. Bull. New York State Mus., 314:1-122.
- Amin, O. M. 1976. Host associations and seasonal occurrence of fleas from southeastern Wisconsin mammals, with observations on morphological variations. J. Med. Entomol., 13:179-192.
- Angelakos, E. T., J. T. Maher, and R. F. Burlington. 1969. Spontaneous cardiac activity at low temperatures in hibernators and non-hibernators: influence of potassium and catecholamines. Fed. Proc., 28:1216-1219.
- Aniskowicz, B. T., and J. Vaillancourt. 1979. Agonistic interactions among wild eastern chipmunks (*Tamias striatus*). Canadian J. Zool., 57:683-690.
- Anthony, A., N. Bachman, and J. L. Frehn. 1963. Histology of digestive organs of the eastern chipmunk, *Tamias striatus*, in relation to reproduction and hibernation. Proc. Pennsylvania Acad. Sci., 37:56-63.
- Anthony, A., D. W. Munro, and A. Stere. 1967. Effect of hibernation on respiration and oxidative phosphorylation in chipmunk liver homogenates. Proc. Pennsylvania Acad. Sci., 40:43-46.
- Baird, S. F. 1857. Report of the Assistant Secretary. Pp. 47-68, in Appendix to the Eleventh Annual Report of the Board of Regents of the Smithsonian Institution . . . for the year 1856.
- 1858. In Mammals. Report of explorations and surveys, to ascertain the most practicable and economical route for a railroad from the Mississippi River to the Pacific Ocean. Washington, D.C., 8(1): 1-757.
- Bangs, O. 1896. Some new mammals from Indian Territory and Missouri. Proc. Biol. Soc. Washington, 10:135-138.
- Bennett, G. F. 1973. Some effects of *Cuterebra emasculator* Fitch (Diptera: Cuterebridae) on the blood and activity of its host, the eastern chipmunk. J. Wildl. Dis., 9:85-93.
- Berry, R. L., et al. 1975. Studies on the epidemiology of California encephalitis in an endemic area in Ohio in 1971. Amer. J. Trop. Med. Hyg., 24:992-998.

- Black, C. C. 1963. A review of the North American Tertiary Sciuridae. *Bull. Mus. Comp. Zool.*, 130:109-248.
- 1972. Holarctic evolution and dispersal of squirrels (Rodentia: Sciuridae). *Evol. Biol.*, 6:305-322.
- Black, J. H., et al. 1977. The eastern chipmunk in central Oklahoma, with validation of Howell's 1929 Red Fork record. *Proc. Oklahoma Acad. Sci.*, 57:163-164.
- Blake, B. H. 1977. The effects of kidney structure and the annual cycle on water requirements in golden-mantled ground squirrels and chipmunks. *Comp. Biochem. Physiol.*, 58A:413-419.
- Bole, B. P., Jr., and P. N. Moulthrop. 1942. The Ohio Recent mammal collection in the Cleveland Museum of Natural History. *Sci. Publ. Cleveland Mus. Nat. Hist.*, 5:83-181.
- Brenner, F. J., and P. D. Lyle. 1975. Effect of previous photoperiodic conditions and visual stimulation on food storage and hibernation in the eastern chipmunk (*Tamias striatus*). *Amer. Midland Nat.*, 93:227-234.
- Brenner, F. J., et al. 1978. Body weight and social interactions as factors in determining dominance in captive eastern chipmunks. *Anim. Behav.*, 26:432-437.
- Bryant, M. D. 1945. Phylogeny of Nearctic Sciuridae. *Amer. Midland Nat.*, 33:257-390.
- Buech, R. R. 1977. Small mammals in a gamma-irradiated northern forest community. Pp. 167-180, in *The Enterprise Wisconsin Radiation Forest. Radioecological Studies* (J. Zavitskovski, ed.). Energy Research and Development Administration, Oak Ridge, Tennessee, TID-26113-P2, 211 pp.
- Burt, W. H. 1940. Territorial behavior and populations of small mammals in southern Michigan. *Misc. Publ. Mus. Zool., Univ. Michigan*, 45:1-58.
- 1960. *Bacula of North American mammals*. *Misc. Publ. Mus. Zool., Univ. Michigan*, 113:1-76.
- Cameron, A. W. 1950. A new chipmunk (*Tamias*) from Ontario and Quebec. *J. Mamm.*, 31:347-348.
- Carey, A. B., W. L. Krinsky, and A. J. Main. 1980. *Ixodes dammini* (Acari: Ixodidae) and associated ixodid ticks in south-central Connecticut, U.S.A. *J. Med. Entomol.*, 17:89-99.
- Catesby, M. 1743. *The natural history of Carolina, Florida, and the Bahama Islands*. . . . Privately printed, London, 2:1-100.
- Clulow, F. V., A. DesMarais, and J. Vaillancourt. 1969. Physiological correlations of population density in the eastern chipmunk (*Tamias striatus*) and the white-footed mouse (*Peromyscus leucopus*). *Canadian J. Zool.*, 47:427-433.
- Dalquest, W. W., E. Roth, and F. Judd. 1969. The mammal fauna of Schulze Cave, Edwards County, Texas. *Bull. Florida State Mus., Biol. Sci.*, 13:205-276.
- DeCoursey, P. J. 1972. LD ratios and the entrainment of circadian activity in a nocturnal and a diurnal rodent. *J. Comp. Physiol.*, 78:221-235.
- Dorney, R. S. 1965. Incidence of botfly larvae (*Cuterebra emasculator*) in the chipmunk (*Tamias striatus*) and red squirrel (*Tamiasciurus hudsonicus*) in northern Wisconsin. *J. Parasitol.*, 51:893-894.
- 1966. Quantitative data on four species of *Eimeria* in eastern chipmunks and red squirrels. *J. Protozool.*, 13:549-550.
- Dudkiewicz, A. B. 1968. Histology of the female reproductive tract in the eastern chipmunk, *Tamias striatus*, and exfoliate cytology of the vagina. Unpubl. M.S. thesis, Univ. Massachusetts, Amherst, 54 pp.
- Duncan, S. 1968. *Eimeria ovata* n. sp. and other coccidia of the eastern chipmunk, *Tamias striatus*, in Massachusetts. *J. Protozool.*, 15:319-320.
- Dunford, C. 1970. Behavioral aspects of spatial organization in the chipmunk, *Tamias striatus*. *Behaviour*, 36:215-231.
- 1972. Summer activity of eastern chipmunks. *J. Mamm.*, 53:176-180.
- Edwards, G. 1751. *A natural history of birds*. . . . College of Physicians, London, 4:158-248. (Although Linnaeus' reference is not specific, the pagination and context indicates this is the edition to which he referred.)
- Ellerman, J. R. 1940. The families and genera of living rodents. *British Mus. Nat. Hist.*, London, 1:1-689.
- Elliott, L. 1978. Social behavior and foraging ecology of the eastern chipmunk (*Tamias striatus*) in the Adirondack Mountains. *Smithsonian Contrib. Zool.*, 265:1-107.
- Ellis, L. S. 1979. Systematics and life-history of the eastern chipmunk, *Tamias striatus*. Unpubl. Ph.D. dissert., Univ. Illinois, Urbana, 325 pp.
- Ellis, L. S., and L. R. Maxson. 1979. Evolution of the chipmunk genera *Eutamias* and *Tamias*. *J. Mamm.*, 60:331-334.
- Elwell, A. S. 1967. Biology of *Anguispira alternata* (Gastropoda: Endodontidae) with special reference to the digenetic trematode *Postharmostomum helices*. *Dissert. Abstr.*, 28:1725B-1726B.
- Engels, W. L. 1951. Winter inactivity of some captive chipmunks (*Tamias s. striatus*) at Chapel Hill, North Carolina. *Ecology*, 32:549-555.
- Fain, A., and K. Hyland. 1970. Notes on the Myocoptidae of North America with description of a new species on the eastern chipmunk, *Tamias striatus* Linnaeus. *J. New York Entomol. Soc.*, 78:80-87.
- Fain, A., et al. 1971. Notes on the hypopial nymphs phoretic on mammals in Canada with description of a new species (Acarina: Sarcoptiformes). *Canadian J. Zool.*, 49:15-18.
- Faleschini, R. J., and B. K. Whitten. 1975. Comparative hypoxic tolerance in the Sciuridae. *Comp. Biochem. Physiol.*, 52A:217-222.
- Fisher, K. E. 1972. Seasonal influences on thyroid histology in the eastern chipmunk (*Tamias striatus*). Unpubl. M.S. thesis, Clarion State College, Clarion, Pennsylvania, 68 pp.
- Forbes, R. B. 1966a. Fall accumulation of fat in chipmunks. *J. Mamm.*, 47:715-716.
- 1966b. Studies of the biology of Minnesotan chipmunks. *Amer. Midland Nat.*, 76:290-308.
- 1967. Some aspects of the water economics of two species of chipmunks. *J. Mamm.*, 48:466-468.
- Freeman, R. S. 1956. Life history studies on *Taenia mustelae* Gmelin, 1790 and the taxonomy of certain taenioid cestodes from Mustelidae. *Canadian J. Zool.*, 34:219-242.
- 1957. Life cycle and morphology of *Paruterina rauschi* n. sp. and *P. candelabraria* (Goeze, 1782) (Cestoda) from owls, and significance of plerocercoids in the order Cyclophyllidae. *Canadian J. Zool.*, 35:349-370.
- 1959. On the taxonomy of the genus *Cladotaenia*, the life histories of *C. globifera* (Batsch, 1786) and *C. circi* Yamaguti, 1935, and a note on distinguishing between the plerocercoids of the genera *Paruterina* and *Cladotaenia*. *Canadian J. Zool.*, 37:317-340.
- Frost, W. W. 1968. Studies on the summer and fall food habits of *Tamias striatus*, the eastern chipmunk, in western Massachusetts. Honors thesis, Dept. Zool., Univ. Massachusetts, Amherst, 101 pp.
- Getty, T. 1979. An observation on *Tamias striatus* reproducing in a tree nest. *J. Mamm.*, 60:636.
- Giles, R. H., Jr. 1970. The ecology of a small forested watershed treated with the insecticide malathion-S³⁵. *Wildl. Monogr.*, 24:5-81.
- Gmelin, J. F. 1788. *Caroli a Linné Systema Naturae* . . . , 13th ed. G. E. Beer, Leipzig, 1:1-500.
- Goodpaster, W. W., and D. F. Hoffmeister. 1968. Notes on Ohioan mammals. *Ohio J. Sci.*, 68:116-117.
- Graefe, G. 1961. Die Aktivitätsperiodik des Flughörnchens, *Glaucomys volans* (Linnaeus 1758) und des Backenhörnchens, *Tamias striatus* (Linnaeus 1758). Ein Beitrag zur Frage der Doppelbesetzung von Gehegen mit dunkel- und lichtaktiven Tierarten. *Z. Tierpsychol.*, 18:84-90.
- Graybill, D. L. 1970. Food resources and ingestion rates of *Tamias striatus* populations. Unpubl. Ph.D. dissert., Univ. Pittsburgh, Pittsburgh, Pennsylvania, 212 pp.
- Guilday, J. E. 1963. Bone refuse from the Oakfield Site, Genesee County, New York. *Pennsylvania Arch.*, 33(1-2):12-15.
- Hall, E. R. 1981. *The mammals of North America*. Second ed. John Wiley and Sons, 1:1-600 + 90.
- Hamilton, W. J., Jr., and J. O. Whitaker, Jr. 1979. Mammals of the eastern United States. 2nd ed. Comstock Publishing Associates, Ithaca, 346 pp.
- Herbig, R. A. 1978. Studies on exploration in the eastern chipmunk. Unpubl. M.A. thesis, Mt. Holyoke College, South Hadley, Massachusetts, 204 pp.
- Hight, M. E., M. Goodman, and W. Prychodko. 1974. Immunological studies of the Sciuridae. *Syst. Zool.*, 23:12-25.
- Hooper, E. T. 1942. Geographic variation in the eastern chipmunk, *Tamias striatus*, in Michigan. *Occas. Pap. Mus. Zool., Univ. Michigan*, 461:1-5.
- Howell, A. H. 1925. Preliminary descriptions of five new chipmunks from North America. *J. Mamm.*, 6:51-54.
- 1929. Revision of the American chipmunks (genus *Tamias* and *Eutamias*). *N. Amer. Fauna*, 52:1-157.

- 1938. Review of the North American ground squirrels with a classification of the North American Sciuridae. N. Amer. Fauna, 56:1-256.
- Ickes, R. A. 1974. Agonistic behavior and the use of space in the eastern chipmunk, *Tamias striatus*. Unpubl. Ph.D. dissert., Univ. Pittsburgh, Pittsburgh, Pennsylvania, 235 pp.
- Illiger, C. 1811. *Prodromus systematis mammalium et avium additis terminis zoographicis utriusque classis*. C. Salfeld, Berlin, 301 pp.
- Jackson, H. H. T. 1961. Mammals of Wisconsin. Univ. Wisconsin Press, Madison, 504 pp.
- Jones, G. S., and H. H. Thomas. 1980. Ticks from mammals from Prince Edward Island, New Brunswick, northern Nova Scotia, and Gaspé Peninsula, Quebec. Canadian J. Zool., 58:1394-1397.
- Kalm, Pehr. 1756. En resa til Norra America . . . Lars Salvii. Stockholm, 2:1-526 + [11 ll]. (Linnaeus' cryptic reference was presumably to this original Swedish edition.)
- Kavanau, J. L. 1969. Influences of light on activity of small mammals. Ecology, 50:548-557.
- Klingener, D. 1968. Rodents of the Mio-Pliocene Norden Bridge local fauna, Nebraska. Amer. Midland Nat., 80:65-74.
- Kodrich, W. R., and C. A. Tryon. 1971. Effect of season on thyroid release of iodine-131 in free ranging eastern chipmunks (*Tamias striatus*). Pp. 260-264, in Radionuclides in ecosystems (D. J. Nelson, ed.). Proc. Third Natl. Symp. Radioecology, Conf-710501-P1, National Technical Information Service, Springfield, Virginia, 1:1-678.
- Kuhl, H. 1820. Beiträge zur Zoologie und vergleichenden Anatomie. Hermann, Frankfurt am Main, 1:1-151.
- Kurtén, B., and E. Anderson. 1980. Pleistocene mammals of North America. Columbia Univ. Press, New York, 442 pp.
- Lankester, M. W., and R. C. Anderson. 1966. Small mammals as paratenic hosts of lungworms. Canadian J. Zool., 44:341-342.
- Layne, J. N. 1954. The os clitoridis of some North American Sciuridae. J. Mamm., 35:357-366.
- 1957. Homing behavior of chipmunks in central New York. J. Mamm., 38:519-520.
- Layne, J. N., and A. H. Benton. 1954. Some speeds of small mammals. J. Mamm., 35:103-104.
- Lindsay, E. H. 1972. Small mammal fossils from the Barstow Formation, California. Univ. Calif. Publ. Geol. Sci., 93:1-104.
- Linnaeus, C. 1754. Mus. Adolphi Friderici Regis . . . Stockholm.
- 1758. Systema Naturae. 10th ed. Laurentii Salvii, Stockholm, 1:1-824.
- 1766. Systema Naturae. 12th ed. Laurentii Salvii, Stockholm, 1:1-1327.
- Long, C. A. 1971. A new subspecies of chipmunk from the Door peninsula, Wisconsin (Mammalia: Rodentia). Proc. Biol. Soc. Washington, 84:201-202.
- Lovejoy, D., and P. J. Gaughan. 1975. Notes on a collection of mammal-fleas from New Hampshire. Entomol. News, 86:145-149.
- Lowery, G. H., Jr. 1943. Check-list of the mammals of Louisiana and adjacent waters. Occas. Pap. Mus. Zool., Louisiana State Univ., 13:213-257.
- 1974. The mammals of Louisiana and its adjacent waters. Louisiana State Univ. Press, Baton Rouge, 565 pp.
- Lyman, C. P., and D. C. Blinks. 1959. The effect of temperature on the isolated hearts of closely related hibernators and non-hibernators. J. Cell Comp. Physiol., 54:53-63.
- Mares, M. A., M. D. Watson, and T. E. Lacher, Jr. 1976. Home range perturbations in *Tamias striatus*. Food supply as a determinant of home range and density. Oecologia, 25:1-12.
- McKinney, T. D., and J. J. Christian. 1970. Incidence and effects of botfly parasitism in the eastern chipmunk. J. Wildl. Dis., 6:140-143.
- Mearns, E. A. 1891. Description of a new subspecies of the eastern chipmunk from the upper Mississippi region, west of the Great Lakes. Bull. Amer. Mus. Nat. Hist., 3:229-233.
- Merriam, C. H. 1886. Description of a new subspecies of the common eastern chipmunk. Amer. Nat., 20:236-242.
- 1897. Notes on the chipmunks of the genus *Eutamias* occurring west of the east base of the Cascade-Sierra system, with descriptions of new forms. Proc. Biol. Soc. Washington, 11:189-212.
- Moore, J. C. 1959. Relationships among living squirrels of the Sciurinae. Bull. Amer. Mus. Nat. Hist., 118:153-206.
- 1961. The spread of existing diurnal squirrels across the Bering and Panamanian Land Bridges. Amer. Mus. Novitates, 2044:1-26.
- Mossman, H. W., J. W. Lawlah, and J. A. Bradley. 1932. The male reproductive tract of the Sciuridae. Amer. J. Anat., 51:89-141.
- Munro, D. W., and A. Anthony. 1965. Respiration and oxidative phosphorylation in liver tissue from hibernating and active chipmunks. Proc. Pennsylvania Acad. Sci., 39:114-120.
- Nadler, C. F., and R. S. Hoffmann. 1970. Chromosomes of some Asian and South American squirrels (Rodentia: Sciuridae). Experientia, 26:1383-1386.
- Nadler, C. F., et al. 1977. Chromosomal evolution in chipmunks with special emphasis on A and B karyotypes of the subgenus *Neotamias*. Amer. Midland Nat., 98:343-353.
- Neal, C. M., and S. I. Lustick. 1975. Energy budget of the eastern chipmunk (*Tamias striatus*): artificial radiation. Comp. Biochem. Physiol., 50A:233-236.
- Neff, W. H. 1968. Gross metabolic response in cold-exposed chipmunks. Proc. Pennsylvania Acad. Sci., 42:81-86.
- Neff, W. H., and A. Anthony. 1963. Seasonal changes in the male reproductive tract of the eastern chipmunk, *Tamias striatus*. Proc. Pennsylvania Acad. Sci., 37:64-70.
- Neumann, R. L. 1967. Metabolism in the eastern chipmunk (*Tamias striatus*) and the southern flying squirrel (*Glaucomys volans*) during the winter and summer. Pp. 64-74, in Mammalian hibernation III (K. C. Fisher et al., eds.). Amer. Elsevier, New York, 535 pp.
- Northcott, T. H., E. Mercer, and E. Menchenton. 1974. The eastern chipmunk *Tamias striatus* in insular Newfoundland. Canadian Field-Nat., 88:86.
- Panuska, J. A. 1959. Weight patterns and hibernation in *Tamias striatus*. J. Mamm., 40:554-566.
- Panuska, J. A., and N. J. Wade. 1956. The burrow of *Tamias striatus*. J. Mamm., 37:23-31.
- Paradiso, J. L. 1963. The type locality of *Tamias striatus fisheri*. J. Mamm., 44:579-580.
- Parsons, M. A. 1962. A survey of the ectoparasites of the wild mammals of New England and New York State. Unpubl. M.S. thesis, Univ. Massachusetts, Amherst, 378 pp.
- Peifer, R. W. 1979. Great blue herons foraging for small mammals. Wilson Bull., 91:630-631.
- Pidduck, E. R., and J. B. Falls. 1973. Reproduction and emergence of juveniles in *Tamias striatus* (Rodentia:Sciuridae) at two localities in Ontario, Canada. J. Mamm., 54:693-707.
- Pivorun, E. B. 1976a. A gradient calorimeter study of normothermic and hibernating eastern chipmunks, *Tamias striatus*. Comp. Biochem. Physiol., 54A:259-261.
- 1976b. A biotelemetry study of the thermoregulatory patterns of *Tamias striatus* and *Eutamias minimus* during hibernation. Comp. Biochem. Physiol., 53A:265-271.
- 1977. Hibernation of a southern subspecies of *Tamias striatus*: thermoregulatory patterns. Amer. Midland Nat., 98:495-499.
- Quay, W. B. 1965. Comparative survey of the sebaceous and sudoriferous glands of the oral lips and angle in rodents. J. Mamm., 46:23-37.
- Rausch, R., and J. D. Tiner. 1948. Studies on the parasitic helminths of the north central states. I. Helminths of Sciuridae. Amer. Midland Nat., 39:728-747.
- Ray, C. E. 1965. A new chipmunk, *Tamias aristus*, from the Pleistocene of Georgia. J. Paleontol., 39:1016-1022.
- Read, C. P. 1949. Studies on North American helminths of the genus *Capillaria* Zeder, 1800 (Nematoda). I. Capillarids from mammals. J. Parasitol., 35:223-230.
- Richardson, J. 1829. Fauna Boreali-Americana. . . The Quadrupeds (Mammalia). John Murray, London, 1:1-300.
- Riedel, D. 1975. Reproduction, structure and host specificity of *Trypanosoma (Herpetosoma) tamiasi* sp. n. from the eastern chipmunk, *Tamias striatus*. J. Protozool., 22:8-17.
- Roberts, E. F. 1976. Patterns of dispersal and other movements in a population of the eastern chipmunk, *Tamias striatus*. Unpubl. Ph.D. dissert., Univ. Massachusetts, Amherst, 110 pp.
- Roth, E. L. 1972. Late Pleistocene mammals from Klein Cave, Kerr County, Texas. Texas J. Sci., 24:75-84.
- Ryan, D. A., and J. S. Larson. 1976. Chipmunks in residential environments. Urban Ecol., 2:173-178.

- Santiago, L., and A. V. Rake. 1973. Rodent DNA reassociation kinetics. *Biochem. Genet.*, 9:275-282.
- Schaffer, R. E. 1971. Calorigenic contribution of brown fat in *Tamias striatus*. *Proc. Indiana Acad. Sci.*, 80:487-488.
- 1973. Seasonal weight changes of *Tamias striatus* in captivity. *Proc. Indiana Acad. Sci.*, 82:434.
- Schaffer, R. E., and R. W. Bullard. 1970. Comparisons of re-warming from natural torpidity and induced hypothermia in chipmunks (*Tamias striatus*) with reference to heart rate and temperature relationships. *Proc. Indiana Acad. Sci.*, 79:476-483.
- Schmid, W. D. 1976. Temperature gradients in the nasal passage of some small mammals. *Comp. Biochem. Physiol.*, 54A:305-308.
- Scott, G. W., and K. C. Fisher. 1972. Hibernation of eastern chipmunks (*Tamias striatus*) maintained under controlled conditions. *Canadian J. Zool.*, 50:95-105.
- Sealander, J. A. 1964. The influence of body size, season, sex, age and other factors upon some blood parameters in small mammals. *J. Mamm.*, 45:598-616.
- Seidel, D. R. 1961. Homing in the eastern chipmunk. *J. Mamm.*, 42:256-257.
- Seton, E. T. 1929. *Lives of game animals*. Doubleday Doran and Co. Inc., Garden City, New York, 4:1-440; 441-949.
- Simpson, S. 1914. The pyramid tract in the red squirrel (*Sciurus hudsonius loquax*) and chipmunk (*Tamias striatus lysteri*). *J. Comp. Neurol.*, 24:137-160.
- Smith, D. A., and L. C. Smith. 1972. Aberrant coloration in Canadian eastern chipmunks, *Tamias striatus*. *Canadian Field-Nat.*, 86:253-257.
- 1975. Oestrus, copulation, and related aspects of reproduction in female eastern chipmunks, *Tamias striatus* (Rodentia: Sciuridae). *Canadian J. Zool.*, 53:756-767.
- Smith, L. C., and D. A. Smith. 1972. Reproductive biology, breeding seasons, and growth of eastern chipmunks, *Tamias striatus* (Rodentia: Sciuridae) in Canada. *Canadian J. Zool.*, 50:1069-1085.
- Snyder, D. P., C. A. Tryon, and D. L. Graybill. 1976. Effect of gamma radiation on range parameters in the eastern chipmunk, *Tamias striatus*. Pp. 354-358, in *Radioecology and energy resources* (Cushing, C. E., ed.). *Proc. Fourth Natl. Symp. Radioecology. Spec. Publ. Ecol. Soc. Amer.*, 1:1-401.
- Stevenson, H. M. 1962. Occurrence and habits of the eastern chipmunk in Florida. *J. Mamm.*, 43:110-111.
- Sutton, J. F. 1972. Additional rodent material from the split-rock local fauna, Miocene of Wyoming. *Occas. Pap. Mus., Texas Tech Univ.*, 4:1-8.
- Symington, R. B. 1907. Frogs catching birds. *Forest and Stream*, 68(16):612.
- Tadkowski, T. M., and K. E. Hyland. 1979. The developmental stages of *Aplodontopus sciuricola* (Astigmata) from *Tamias striatus* L. (Sciuridae) in North America. Pp. 321-326, in *Proc. 4th Int. Cong. Acarology, 1974* (E. Piff, ed.). *Hungarian Acad. Sci., Budapest*, 752 pp.
- Thibault, P. 1969. Activité estivale de petits mammifères du Québec. *Canadian J. Zool.*, 47:817-828.
- Thomas, K. R. 1974. Burrow systems of the eastern chipmunk (*Tamias striatus pipilans* Lowery) in Louisiana. *J. Mamm.*, 55:454-459.
- Thompson, J. N., Jr. 1971. Chromosomes of Oklahoma mammals. I. Three species of squirrels. *Proc. Oklahoma Acad. Sci.*, 51:79-80.
- Tiner, J. D. 1950. Two new species of *Trichuris* from North America with redescription of *Trichuris opaca* and *Trichuris leporis* (Nematoda: Aphasmidia). *J. Parasitol.*, 36:350-355.
- Trouessart, E.-L. 1880. Catalogue des mammifères vivants et fossiles. *Ordre des Rongeurs. Bull. Soc. Etudes Sci. Angers*, 10:58-212.
- Tryon, C. A., and D. P. Snyder. 1973a. Biology of the eastern chipmunk, *Tamias striatus*: life tables, age distributions, and trends in population numbers. *J. Mamm.*, 54:145-168.
- 1973b. The effect of exposure to 200 and 400 r of ionizing radiation on the survivorship curves of the eastern chipmunk (*Tamias striatus*) under natural conditions. Pp. 1037-1041, in *Radionuclides in ecosystems* (D. J. Nelson, ed.). *Proc. Third Natl. Symp. Radioecology, Conf-710501-P2, National Technical Information Service, Springfield, Virginia*, 2:679-1268.
- Van Cleave, H. J. 1953. *Acanthocephala of North American Mammals*. *Illinois Biol. Monogr.*, 23(1-2):1-179.
- Wade, O., and P. T. Gilbert. 1940. The baculum of some Sciuridae and its significance in determining relationships. *J. Mamm.*, 21:52-63.
- Wang, L. C.-H., and J. W. Hudson. 1971. Temperature regulation in normothermic and hibernating eastern chipmunk, *Tamias striatus*. *Comp. Biochem. Physiol.*, 38A:59-90.
- West, G. G. II. 1975. Factors affecting home range and range length of the eastern chipmunk (*Tamias striatus*) in northwestern Pennsylvania. Unpubl. Ph.D. dissert., Univ. Pittsburgh, Pittsburgh, Pennsylvania, 254 pp.
- White, J. A. 1953. Genera and subgenera of chipmunks. *Univ. Kansas Publ., Mus. Nat. Hist.*, 5:543-561.
- Wolfe, J. L. 1966. Agonistic behavior and dominance relationships of the eastern chipmunk, *Tamias striatus*. *Amer. Midland Nat.*, 76:190-200.
- 1969. Observations on alertness and exploratory behavior in the eastern chipmunk. *Amer. Midland Nat.*, 81:249-253.
- Woods, C. A., and E. B. Howland. 1977. The skin musculature of hystricognath and other selected rodents. *Anat. Histol. Embryol.*, 6:240-264.
- Woodward, A. E., and J. M. Condren. 1945. Physiological studies on hibernation in the chipmunk. *Physiol. Zool.*, 18:162-167.
- Wrazen, J. A., and G. E. Svendsen. 1978. Feeding ecology of a population of eastern chipmunks (*Tamias striatus*) in southeast Ohio. *Amer. Midland Nat.*, 100:190-201.
- Wright, R. E., and G. R. DeFoliart. 1970. Associations of Wisconsin mosquitoes and woodland vertebrate hosts. *Ann. Entomol. Soc. Amer.*, 63:777-786.
- Yahner, R. H. 1975. The adaptive significance of scatter hoarding in the eastern chipmunk. *Ohio J. Sci.*, 75:176-177.
- 1977. Activity lull of *Tamias striatus* during the summer in southeast Ohio. *Ohio J. Sci.*, 77:143-145.
- 1978a. Weight gain of post-emergent juvenile *Tamias striatus*. *J. Mamm.*, 59:196-197.
- 1978b. Burrow system and home range use by eastern chipmunks, *Tamias striatus*: ecological and behavioral considerations. *J. Mamm.*, 59:324-329.
- 1978c. The adaptive nature of the social system and behavior in the eastern chipmunk, *Tamias striatus*. *Behav. Ecol. Sociobiol.*, 3:397-427.
- 1978d. The sequential organization of behavior in *Tamias striatus*. *Behav. Biol.*, 24:229-243.
- Yahner, R. H., B. L. Allen, and W. J. Peterson. 1979. Dorsal and anal glands in the eastern chipmunk, *Tamias striatus*. *Ohio J. Sci.*, 79:40-43.
- Yerger, R. W. 1953. Home range, territoriality, and populations of the chipmunk in central New York. *J. Mamm.*, 34:448-458.
- 1955. Life history notes on the eastern chipmunk, *Tamias striatus lysteri* (Richardson), in central New York. *Amer. Midland Nat.*, 53:312-323.

Editors of this account were DANIEL F. WILLIAMS and SYDNEY ANDERSON. Managing editor was TIMOTHY E. LAWLOR.

D. P. SNYDER, DEPARTMENT OF ZOOLOGY, UNIVERSITY OF MASSACHUSETTS, AMHERST 01003.